

McGill/96-17
 hep-ph/9606232
 May, 1996

R_b PROBLEM: LOOP CONTRIBUTIONS AND SUPERSYMMETRY

J.M. CLINE

McGill University, 3600 University St., Montréal, Québec H3A 2T8, Canada

I summarize a part of the work done with Bamert, Burgess, London and Nardi,¹ in which we find simplified expressions for the possible one-loop contributions to the R_b parameter from new virtual particles, including supersymmetric ones. Our expressions make it easier to identify which models and choices of parameters are best able to solve the R_b problem.

The evidence for a discrepancy between the standard model prediction for R_b , the ratio of partial decay widths for $Z \rightarrow b\bar{b}$ to $Z \rightarrow \text{hadrons}$, has been described by P. Bamert.² In this talk I will recapitulate our investigation¹ of the possibility of resolving the R_b problem using new particles in the loops of the diagrams of Figure 1, which contribute to the $Z \rightarrow b\bar{b}$ width. For example, possible combinations of particles in the loop could be an extra Higgs boson and the top quark, a squark and a chargino, or the usual Higgs with a 4th generation quark. Many different extensions of the standard model have been suggested for solving the R_b problem.³ However, it would be nice if it was not necessary to compute the corrections to R_b anew for every such possibility. Our goal in this study is to identify what are the new physics ingredients needed to increase the value of R_b through one-loop effects, in as model-independent a way as possible.

1 Methodology

To achieve our goal, it is very useful to make three simplifications.

(1) Ignore corrections to the $Zb_R\bar{b}_R$ vertex. The left- and right-handed b quarks couple to the Z boson through the interaction

$$\frac{1}{2}\bar{b}\gamma^\mu (g_L^b(1 - \gamma_5) + g_R^b(1 + \gamma_5)) b Z_\mu. \quad (1)$$




Figure 1: Loop corrections to the $Z \rightarrow b\bar{b}$ decay width.

As already discussed,² the tree-level value of g_R^b is so small that we would need to increase it by 100% if we wanted to solve the R_b problem just by changing g_R^b . Such a large shift would be quite difficult to achieve from loop corrections for any reasonable-size coupling constants. On the other hand, to solve the R_b problem by increasing g_L^b is possible at one loop, as is demonstrated by the fact that the absence of the top quark loop correction to R_b in the standard model would be enough to remove the discrepancy.

(2) Set the external momentum q of the Z boson to $q^2 = 0$ rather than its actual value of $q^2 = m_Z^2$. This makes the formulas for the loop diagrams much more tractable, and typically introduces only a small error, as I will show below.

(3) Ignore vacuum polarization corrections, as in figure 2. At one-loop accuracy we are justified in doing so, because they largely cancel in the ratio which defines R_b . Let us denote the relative corrections to the hadronic widths due to the vacuum polarization diagrams of figure 2 by ϵ_{vp} , and the absolute corrections from figure 1 to the partial widths into $b\bar{b}$ and hadrons, respectively, by ϵ_b and ϵ_h :

$$R_b \equiv \frac{\Gamma_b}{\Gamma_{\text{hadrons}}} = \frac{\Gamma_b^0(1 + \epsilon_{vp}) + \epsilon_b}{\Gamma_h^0(1 + \epsilon_{vp}) + \epsilon_h} = \frac{\Gamma_b^0 + \epsilon_b + \mathcal{O}(\epsilon_b\epsilon_{vp})}{\Gamma_h^0 + \epsilon_h + \mathcal{O}(\epsilon_b\epsilon_{vp})}, \quad (2)$$

where Γ_i^0 is the tree-level value. Because ϵ_{vp} is a universal correction for all flavors of quarks, it cancels in the ratio, to one loop accuracy.




Figure 2: Vacuum polarization corrections to the $Z \rightarrow b\bar{b}$ decay width, which we ignore.

This last simplification is helpful not only because it reduces the number of diagrams that must be calculated, but also because the ones that remain (figure 1) have nice properties. Thanks to the Ward identity, the diagrams 1a and 1b are related to those of 1c and 1d in such a way that the sum of the four is finite and gauge invariant. Let us recall how this works. If p' and p are the initial and final momenta of the b quark, respectively, then the Ward identity is

$$\begin{aligned} (p - p')^\mu \Gamma_\mu &= g (S_F^{-1}(p) - S_F^{-1}(p')) ; \\ (g + \delta g)(\not{p} - \not{p}') &= g(1 + \delta Z)(\not{p} - \not{p}'). \end{aligned} \quad (3)$$

Here Γ_μ is the 1PI contribution to the vertex correction, which can be written as $(g + \delta g)\gamma_\mu$, in terms of the tree-level coupling g and its one-loop shift δg , and δZ is the wave function renormalization. Thus diagrams 1a and 1b are associated with δg and similarly 1c and 1d with δZ . Although the Ward identity is derived for an unbroken $U(1)$ symmetry, the divergent parts of the diagrams are the same for the Z boson as for the photon, so the sum of the diagrams is finite. Furthermore, we shall see that in our approximation of $q^2 = 0$ (point 2 above), the parts of the diagrams proportional to $\sin^2\theta_W$ combine to give zero. Again, these are the parts proportional to the photon coupling, and the neglect of the Z mass makes the corresponding $U(1)$ symmetry appear to be unbroken.

2 Results

Before presenting the results for the loop diagrams, here are our conventions for the couplings of the new fermion (f) and scalar (ϕ) in the loop to the Z boson and to each other:

$$\begin{aligned} \mathcal{L}_{\text{gauge}} &= \frac{e}{\sin 2\theta_W} \left[\bar{f} \gamma^\mu \left(g_L^f (1 - \gamma_5) + g_R^f (1 + \gamma_5) \right) f + 2i g^\phi \phi^\dagger \overleftrightarrow{\partial}_\mu \phi \right] Z_\mu ; \\ \mathcal{L}_{\text{Yukawa}} &= \frac{y}{2} \bar{b} \phi (1 + \gamma_5) f + \text{h.c.} \end{aligned} \quad (4)$$

Notice that we have not bothered to write a Yukawa coupling to b_R , the right-handed b quark, since we are only interested in generating corrections to g_L^b . Moreover gauge invariance demands the relation $g_L^b = g^\phi + g_L^f$. The gauge couplings are related to the third component of weak isospin and the electric charge in the usual way,

$$g = (T_3 - Q \sin^2\theta_W). \quad (5)$$




Figure 3: The function $\mathcal{F}$ of the fermion to scalar mass ratio from eq. (6).

It is these $\sin^2\theta_W$'s appearing in the gauge couplings that cancel out of all the results in the limit of vanishing m_Z .

The one-loop shift in g_L^b due to the diagrams of figure 1 now has a very simple form,

$$\delta g_L^b = \frac{y^2 n_c}{16\pi^2} (g_L^f - g_R^f) \mathcal{F}(m_f^2/m_\phi^2); \quad \mathcal{F}(r) = \frac{r}{r-1} - \frac{r \ln r}{(r-1)^2}, \quad (6)$$

where n_c is a color factor (*e.g.*, $n_c = 2$ if the particles in the loop are color triplets) and $\mathcal{F}$ is a positive function of the ratio of the fermion to the scalar mass, with $0 \leq \mathcal{F} \leq 1$, as shown in figure 3. Although the gauge coupling of the scalar, g^ϕ , appears in the calculation of δg_L^b , it can be eliminated in favor of the chiral couplings of the fermion, due to gauge invariance. As advertised, the $\sin^2\theta_W$ -dependence has disappeared from the result, and $(g_L^f - g_R^f)$ depends only on the isospins of the two chiral fermions in the loop.

We can immediately discern from eq. (6) what is needed to increase R_b : the combination $(g_L^f - g_R^f)$ must have the same sign (negative) as g_L^b itself. Thus two-Higgs doublet models will not work, to the extent that the top quark Yukawa coupling dominates in the loop over that of the bottom, since $(g_L^t - g_R^t)$ has the opposite sign to g_L^b . This computation also shows why the large top quark mass exacerbates the R_b problem in the standard model itself.

However, one might be concerned about our neglect of the mass of the Z boson. It is straightforward to make an expansion in powers of m_Z^2 over the other masses to see how big the error is. For simplicity consider the special

case where the fermion and the scalar in the loop are degenerate, $m_f = m_\phi$. Then to first order in m_Z^2/m_f^2 , eq. (6) becomes

$$\delta g_L^b = \frac{y^2 n_c}{32\pi^2} \left(g_L^f - g_R^f + \frac{m_Z^2}{12m_f^2} (g_L^b + g_L^f + g_R^f) \right). \quad (7)$$

Thus even if the new particles in the loop were as light as the Z boson, the correction due to the nonvanishing Z mass would still only be $\sim 10\%$ of the leading order result.

3 Nondiagonal Z -couplings and supersymmetry

So far we assumed that the new particles in the loop consisted of a single fermion and scalar, but it is possible that there are several of each, and that their couplings to the Z boson, although diagonal in the flavor basis, are no longer diagonal in the basis of mass eigenstates. For example two flavors of fermions a and b would have the chiral rotation matrices $U_{L,R}$ defined by

$$\begin{pmatrix} f_{L,R}^a \\ f_{L,R}^b \end{pmatrix} = \begin{pmatrix} \cos \theta_{L,R} & \sin \theta_{L,R} \\ -\sin \theta_{L,R} & \cos \theta_{L,R} \end{pmatrix} \begin{pmatrix} f_{L,R}^1 \\ f_{L,R}^2 \end{pmatrix} \quad (8)$$

and similarly for the bosons. This is just what happens in supersymmetric models, where the fermions are the higgsino $\tilde{h}_2^-$ which couples to the bottom quark and top squark with the top quark Yukawa coupling y_t , and the Wino, $\tilde{W}^-$. Their mass matrix, in the basis $(\tilde{h}_2^-, \tilde{W}^-)$, is

$$m_{\text{chargino}}^2 = \begin{pmatrix} \mu & g v_2 \\ g v_1 & M_2 \end{pmatrix} \quad (9)$$

which is generally nonsymmetric, so that it is diagonalized by a similarity transformation with $U_L \neq U_R$. Likewise there are two bosons, the right- and left-handed top squarks $\tilde{t}_{R,L}$ with mass matrix

$$m_{\text{squark}}^2 = \begin{pmatrix} m_U^2 + m_t^2 + \mathcal{O}(m_W^2) & m_t(A_t \sin \beta + \mu \cos \beta) \\ m_t(A_t \sin \beta + \mu \cos \beta) & m_Q^2 + m_t^2 + \mathcal{O}(m_W^2) \end{pmatrix}. \quad (10)$$

In the flavor basis, the gauge coupling constants of the previous section become matrices,

$$g_L^f = g_R^f = \begin{pmatrix} -\frac{1}{2} & 0 \\ 0 & -1 \end{pmatrix}, \quad g^\phi = \begin{pmatrix} 0 & 0 \\ 0 & \frac{1}{2} \end{pmatrix}, \quad (11)$$

where we ignored the $\sin^2 \theta_W$ parts as justified above.

The generalized expression for δg_L^b corresponding to eq. (6) is much more complicated when we have multiple particles with mixing in the loop.^a It can be written in terms of some elementary double integrals,

$$\begin{aligned} \delta g_L^b = & \frac{y^2 n_c}{32\pi^2} \int_0^1 du \int_{-u}^u dv \sum_{ijkl} U_{1i}^\phi U_{1j}^{\phi*} U_{1k}^R U_{1l}^{R*} \left\{ g_L^b \delta_{ij} \delta_{kl} \ln(u m_{\phi_i}^2 + (1-u) m_{f_l}^2) \right. \\ & - (U_R^\dagger g_R^f U_R)_{kl} \delta_{ij} \left(1 + \ln \left((1-u) m_{\phi_i}^2 + \frac{u}{2} (m_{f_k}^2 + m_{f_l}^2) - \frac{v}{2} (m_{f_k}^2 - m_{f_l}^2) \right) \right) \\ & + \delta_{ij} m_{f_k} (U_L^\dagger g_L^f U_L)_{kl} m_{f_l} \left((1-u) m_{\phi_i}^2 + \frac{u}{2} (m_{f_k}^2 + m_{f_l}^2) - \frac{v}{2} (m_{f_k}^2 - m_{f_l}^2) \right)^{-1} \\ & \left. - \delta_{kl} (U_\phi^\dagger g_\phi^f U_\phi)_{ij} \ln \left(\frac{u}{2} (m_{\phi_i}^2 + m_{\phi_j}^2) + (1-u) m_{f_k}^2 - \frac{v}{2} (m_{\phi_i}^2 - m_{\phi_j}^2) \right) \right\}. \quad (12) \end{aligned}$$

Although it is hard to glean much from this formula by inspection, there are three enlightening special cases where the result again becomes simple.

(1) No mixing. If $\mu, M_2 \gg m_W$ in the chargino mass matrix and $m_t A_t, m_t \mu \ll m_i^2 + m_{Q,U}^2$ in the squark mass matrix there would be small mixing angles in both sectors. Then the simple formula (6) applies, using the 1-1 elements of g_L^f and g_R^f , since these are the states that are defined to couple to the left-handed b quark in the absence of mixing. In the SUSY case we see immediately that δg_L^b is zero in this limit, since $g_L^f - g_R^f = 0$.

(2) Mixing, but $m_{\phi_1} = m_{\phi_2}$ and $m_{f_1} = m_{f_2}$. (Degenerate masses imply no mixing in the scalar sector, but there is generally still mixing of the fermions because the asymmetry of the Dirac mass matrix.) Then eq. (6) again applies, but with the replacement

$$g_L^f - g_R^f \rightarrow \left(U_R U_L^\dagger g_L^f U_L U_R^\dagger - g_R^f \right)_{11}. \quad (13)$$

With maximal mixing so that $U_R U_L = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, we would have $U_R U_L^\dagger g_L^f U_L U_R^\dagger = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -\frac{1}{2} & 0 \\ 0 & -1 \end{pmatrix}$ so that

$$\left(U_R U_L^\dagger g_L^f U_L U_R^\dagger - g_R^f \right)_{11} = -1 + \frac{1}{2} = -\frac{1}{2}, \quad (14)$$

which has the correct sign to increase R_b ! Thus we see that large mixing in the chargino sector is desirable for boosting R_b . The condition can be written in terms of the mixing angles as

$$\begin{aligned} U_R U_L^\dagger &= \begin{pmatrix} \cos \theta_R & \sin \theta_R \\ -\sin \theta_R & \cos \theta_R \end{pmatrix} \begin{pmatrix} \cos \theta_L & -\sin \theta_L \\ \sin \theta_L & \cos \theta_L \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ \implies \tan \theta_R \tan \theta_L &= -1 \end{aligned} \quad (15)$$

^aFor detailed formulas in the case of the minimal supersymmetric standard model, see ref. 4.

It is straightforward to work out the consequences of this restriction for the chargino mass matrix, eq. (9). One finds that light charginos are required, such that $m_{\chi_1} + m_{\chi_2} \cong 2m_W$. Furthermore $\tan\beta$ (the ratio of the two Higgs VEV's $\langle H_2 \rangle / \langle H_1 \rangle$) is restricted to lie in the range $m_{\chi_1}/m_{\chi_2} \leq \tan\beta \leq m_{\chi_2}/m_{\chi_1}$. Given the restriction on the sum of the chargino masses, their ratio can never be very large or very small without having one of them be lighter than the experimental limit, and thus it is necessary to have $\tan\beta \sim 1$. In addition, the shift in R_b is proportional to the function $\mathcal{F}$ of eq. (6), which will suppress the result if $\tilde{t}_R$ is much heavier than the charginos. Thus we also see that a relatively light right-handed top squark is needed.

(3) Heavy scalars. In the limit that $m_{\phi_1}, m_{\phi_2} \gg m_{f_1}, m_{f_2}$ the expression for δg_L^b again simplifies:

$$\delta g_L^b = \frac{y^2 n_c}{64\pi^2} \sin^2 2\theta_\phi \left(g_{22}^\phi - g_{11}^\phi \right) \mathcal{F}'(m_{\phi_1}^2/m_{\phi_2}^2) \quad (16)$$

where θ_ϕ is the scalar mixing angle and $\mathcal{F}'$ is another nonnegative function. All of the factors in this equation are thus positive with the possible exception of $g_{22}^\phi - g_{11}^\phi$, hence the sign of the latter determines the sign of δg_L^b . In the SUSY case $g_{22}^\phi - g_{11}^\phi = \frac{1}{2}$, which has the wrong sign. Thus one cannot have both top squarks much heavier than the charginos, in agreement with the conclusion of the previous paragraph.

Finally I show some quantitative predictions for δg_L^b without going to the any of the three limits just discussed. To completely account for the R_b discrepancy, one wants a shift of $\delta g_L^b = -0.0063$. Figure 4 shows how δg_L^b depends on various pairs of parameters in supersymmetric extensions of the standard model. In figure 4(a) we vary the right-handed top squark masses and the squark mixing angle, keeping all other masses fixed and degenerate, but with the wrong sign, $\tan\theta_L \tan\theta_R = +1$, relative to condition (15). One sees that the sign of δg_L^b is opposite to that needed for increasing R_b , in accordance with the limiting case (2) above. Figure 4(b) shows the same thing except with condition (15) fulfilled, so that δg_L^b indeed has the desired sign, and also a large enough magnitude if $\tilde{t}_R$ is sufficiently light^b or has large enough mixing. Figure 4(c) shows that large negative values of δg_L^b occur for values of the chargino mixing angles in accordance with (15). However we set $\delta g_L^b = 0$ in the figure for any parameters giving theoretically disfavored values of $\tan\beta < 1$, which is why there are no large values of δg_L^b when $\tan\theta_R$ becomes large, as eq. (15) taken by itself would have predicted. Figure 4(a) also shows the preference for $\tan\theta_L \tan\theta_R = -1$, since here we have set $\tan\theta_L = -1$ and allowed $\tan\theta_R$ to

^bSince we set $m_Z = 0$, the scale of mass is set by that of the charginos. Thus a mass of 1 on figure 4 corresponds to the squarks and charginos all being degenerate with each other.




Figure 4: Dependence of δg_L^b on various parameters in supersymmetric models.

vary. The largest shifts in δg_L^b occur when $\tan \theta_R = +1$. In addition, the mass of the mostly-Wino component of the chargino is allowed to vary, showing that the shift in δg_L^b is sufficiently large as long as the former is not much lighter than the other chargino.

4 Conclusions

In summary, we have found simple, analytic expressions for the shift in the coupling of the left-handed b quark to the Z boson, induced by one-loop diagrams due to new particles beyond the standard model. These expressions make it easy to understand which models are likely to be able to resolve the R_b problem, and to identify the favored regions of parameter space. For example our formulas immediately show that in the simplest supersymmetric extensions of the standard model, one needs $\tan \beta \sim 1$, and light charginos and right-handed top squarks, with masses near m_W . This agrees with and, we hope, illuminates the results of numerous more detailed studies.⁵ Moreover our results make it easier to see how to construct models that can do the job the more effectively, should the need arise. For example, if the favored regions of SUSY parameter

space mentioned above should be ruled out, one could loosen the constraints on the chargino masses by providing new sources of off-diagonal terms in the chargino mass matrix that would be larger than the weak scale. This would allow the charginos to have large mixing angles without requiring them to be light, and give more freedom in the choice of $\tan\beta$.

Acknowledgments

Our work received financial support from NSERC of Canada and FCAR du Québec.

References

1. P. Bamert, C.P. Burgess, J.M. Cline, D. London and E. Nardi, preprint McGill-96-04 (1996), hep-ph/9602438.
2. P. Bamert, in these proceedings, hep-ph-9606227; preprint McGill-95-64 (1995), hep-ph/9512445.
3. J.T. Liu and D. Ng, *Phys. Lett. B***342**, 262 (1995), hep-ph/9408375;
D. Garcia and J. Sola, *Phys. Lett. B***354**, 335 (1995), hep-ph/9502317;
J.K. Kim, Y.G. Kim, J.S. Lee and K.Y. Lee, *Phys. Rev. D***53**, 1712 (1996); hep-ph/9509319;
E.H. Simmons, R.S. Chivukula and J. Terning, preprint BU-HEP-95-27 (1995), hep-ph/9509392;
E. Ma, *Phys. Rev. D***53**, 2276 (1996), hep-ph/9510289
G. Bhattacharyya, G.C. Branco and W.-S. Hou, to appear in *Phys. Rev. D*, hep-ph/9512239 ;
T. Yoshikawa, Hiroshima preprint HUPD-9528 (1995) hep-ph/9512251;
C.-H.V. Chang, D. Chang and W.-Y. Keung, National Tsing Hua U. preprint NHCU-HEP-96-1, hep-ph/9601326;
G. Altarelli, N. Di Bartolomeo, F. Feruglio, R. Gatto and M.L. Mangano, preprint CERN-TH-96-20 (1996), hep-ph/9601324;
K.S. Babu, C. Kolda and J. March-Russell, preprint IASSNS-HEP-96-20 (1996), hep-ph/9603212;
D. Atwood, L. Reina and A. Soni, preprint CEBAF-TH-96-01 (1996), hep-ph/9603210;
D. Comelli and J.P. Silva, to appear in *Phys. Rev. D* (1996), hep-ph/9603221;
J.S. Lee and J.K. Kim, Korea Inst. Sci. preprint KAIST-CHEP-96-03 (1996), hep-ph/9603225;
P.H. Chankowski and S. Pokorski, preprint IFT-96-6 (1996), hep-

- ph/9603310;
U. Mahanta, Mehta Research Inst. preprint MRI-PHY-15-96 (1996), hep-ph/9604365.
4. M. Boulware and D. Finnell, *Phys. Rev. D***44**, 2054 (1991).
 5. X. Wang, J.L. Lopez and D.V. Nanopoulos, CERN-TH-7553-95 (1995), hep-ph/9501258; *Phys. Rev. D***52**, 4116 (1995), hep-ph/9506217;
J.D. Wells and G.L. Kane, *Phys. Rev. Lett.* **76**, 869 (1996), hep-ph/9510372;
J.L. Feng, N. Polonsky and S. Thomas, *Phys. Lett. B***370**, 1996 (,) hep-ph/9511324;
Y. Yamada, K. Hagiwara and S. Matsumoto, preprint KEK-TH-459 (1995), hep-ph/9512227;
J. Ellis, J.L. Lopez and D. Nanopoulos, preprint CERN-TH-95-314 (1995), hep-ph/9512288;
E.H. Simmons and Y. Su, preprint BUHEP-96-4, hep-ph/9602267;
W. Hollik, preprint hep-ph/9603329 (1996).